

April 7 1997
TAUP 2416/97

HARD INCLUSIVE PRODUCTION AND A VIOLATION OF THE FACTORIZATION THEOREM

E. G O T S M A N ^{a)1)}, E. L E V I N ^{a)b)2)} and U. M A O R ^{a)3)}

^{a)}*School of Physics and Astronomy
Raymond and Beverly Sackler Faculty of Exact Science
Tel Aviv University, Tel Aviv, 69978, ISRAEL*

^{b)}*Theory Department, Petersburg Nuclear Physics Institute
188350, Gatchina, St. Petersburg, RUSSIA*

Abstract:

A new mechanism for hard inclusive production, which leads to a violation of the factorization theorem, is suggested. The mechanism is illustrated by a detailed discussion of Higgs meson production in high energy deuteron-deuteron scattering. Numerical estimates for the effect are given for high energy hadron (nucleus) scattering.

¹⁾ Email: gotsman@post.tau.ac.il .

²⁾ Email: leving@post.tau.ac.il .

³⁾ Email: maor@post.tau.ac.il .

In this letter we consider a new mechanism for hard inclusive production, which violates the factorization theorem [1]. We illustrate this mechanism for the case of inclusive heavy Higgs production in deuteron-deuteron high energy scattering. The suggested mechanism is present in any central hard inclusive production, such as the high p_T inclusive production of mini jets and jets.

The usual description of inclusive production of a Higgs meson in a deuteron-deuteron interaction, is given by a simple Mueller diagram [2] shown in Fig.1a. Our calculation is based on the impulse approximation in which the production mechanism is initiated by a primary single nucleon-nucleon collision. Assuming the factorization theorem, the cross section of interest is given by

$$\sigma(Higgs) = \int dx_1 dx_2 F_D^p(x_1, M^2) F_D^p(x_2, M^2) \sigma(hard) , \quad (1)$$

where $F_D^p(x, M^2)$ denotes the parton distribution within the deuteron, and in the impulse approximation $F_D^p = 2F_N^p$. $\sigma(hard)$ denotes the cross section for Higgs meson production in the parton - parton collision.

Since the deuteron is a two nucleon bound state, we may also consider a production mechanism in which two nucleon-nucleon interactions are responsible for the Higgs production. These are shown in Fig.1b which, after squaring, gives the Mueller diagrams of Fig.1c. However, to obtain the full answer a more careful treatment is required in which we sum over all contributions to the inclusive production, according to the AGK cutting rules [3]. This procedure leads to a cancelation of the contributions shown in Fig.1c, so that the net inclusive cross section is given by Fig.1a.

The above consideration fails to include the interference diagram shown in Fig.1d. We claim that this contribution survives the cancelations implied by the AGK cutting rules, and this results in a violation of the factorization theorem. In this note we calculate this contribution and assess the consequences in detail. We wish to remind the reader that the application of the AGK cutting rules are different when calculating inelastic and inclusive processes. This is shown in Fig.2 for nucleon-nucleon scattering.

The present paper continues the approach utilized in many cascade models for inclusive production in nucleus-nucleus interactions (see, for example, [4]). Recently, this mechanism has been revived by Ryskin [5], who pointed out its importance for nucleus-nucleus collisions where its contribution is proportional to $(A_1 A_2)^{\frac{4}{3}}$, whereas using the factorization theorem one obtains only an $(A_1 A_2)$ dependence. The result obtained by Ryskin is evident from Fig.1d as the contribution of this diagram is proportional to $(A_1^2 A_2^2)$ - the number of two nucleon-nucleon collisions - divided by $\pi R_{A_1}^2 \pi R_{A_2}^2$ - which comes from the integration over q_1 and q_2 (the momentum transfers in Fig.1d). Our paper differs from

Refs. [4] [5], in as much as we have included explicitly the relevant contributions and cancelations implied by the AGK cutting rules. For a deuteron-deuteron collision, we find that the interference diagram resulting from the two nucleon interaction (Fig.1d) has to be added to the one nucleon interaction of Fig.1a. Checking the numerics, we obtain a significant addition to the cross section for central mass production, which is in the few GeV range. This is relevant for estimates of the minijet cross section.

In Fig.3 we show the diagrams illustrating the new mechanism for a centrally produced Higgs meson. Fig.3a shows the process where the Higgs, which is produced centrally, is separated by two large rapidity gaps (LRG) from the small final state multiplicities which occur on the edges of the rapidity plot. We define this process as a double Pomeron exchange reaction, and denote its contribution as $\sigma_2^{(0)}$. The contribution of Fig.3b is denoted by $\sigma_2^{(1)}$. This is a mixed diagram, where the Higgs which is produced in a double Pomeron process, is superimposed on a normal uniform rapidity distribution typical of an inelastic nucleon-nucleon reaction. Finally, Fig.3c describes Higgs production as part of the nucleon-nucleon background to the rapidity distribution. We denote this contribution by $\sigma_2^{(2)}$. To obtain our final result we need to sum over all three of the above contributions, noting that these are not necessarily positive.

The nucleon-nucleon (NN) amplitude for Higgs production via double Pomeron exchange, has been calculated [6], to be

$$A_H = A(NN \rightarrow N + (LRG) + H + (LRG) + N) = 2g_H A_P, \quad (2)$$

g_H is the the vertex of the hard parton - parton $\rightarrow$ Higgs process and A_P is the Pomeron exchange amplitude. Fig.3d illustrates graphically the source of the factor 2 (in Eq. (2)), which plays an important role in our calculation. The second ingredient in our calculation is the amplitude shown in Fig.3e. This amplitude has no analog in the case of a single nucleon-nucleon interaction, and it depicts the cut in the diagram shown in Fig.3b. We note that this diagram is equal to ImA_H , unlike the case for inelastic nucleon-nucleon cross section where $\sigma_{in} = 2ImA_P$. The above result follows from the unitarity constraint [3].

Recalling that the integration over the longitudinal components of the vector q_μ in Fig.3b results in a negative sign for the interference diagram (see Ref. [3] for details), we can easily calculate all diagrams contributing to the inclusive Higgs meson production. Indeed, calculating $\sigma_2^{(0)}$, the sum of the diagrams in Fig.3a, and using Eq. (2), one has

$$\sigma_2^{(0)} = 8g_H^2 \{ (ReA_P)^2 + (ImA_P)^2 \}. \quad (3)$$

For the sum of diagrams in Fig.2b we have

$$\sigma_2^{(1)} = -16g_H^2 (ImA_P)^2. \quad (4)$$

Finally, for $\sigma_2^{(2)}$, shown in Fig.2c, we obtain

$$\sigma_2^{(2)} = 8 g_H^2 (Im A_P)^2 . \quad (5)$$

The relation between the different contributions is

$$\sigma_2^{(0)} : \sigma_2^{(1)} : \sigma_2^{(2)} = (1 + \rho^2) : -2 : 1 , \quad (6)$$

where $\rho = \frac{Re A_P}{Im A_P}$. This should be compared with the AGK relation[3] for the inelastic cross sections

$$\sigma_{in}^{(0)} : \sigma_{in}^{(1)} : \sigma_{in}^{(2)} = (1 + \rho^2) : -4 : 2 . \quad (7)$$

Summing all contributions we obtain an additional term in the cross section compare to the one given in Eq.(1)

$$\sigma(D + D \rightarrow H + X) = 8 g_H^2 (Re A_P)^2 . \quad (8)$$

We would like to draw attention to the novel fact, that the contribution of the new mechanism is proportional to the real part of the amplitude. For a soft Pomeron with $\alpha_P(0) = 1$, the contribution of Eq.(8) vanishes, and we recover the factorization theorem [1], i.e. an exact cancellation of the diagrams shown in Fig.1c.

The detailed kinematics of a double Pomeron Higgs meson production, as well as its amplitude, have been discussed in Ref.[6]. The most important kinematical feature is that at high energy and for a Higgs meson in the central rapidity region: $y \gg 1$, and $Y - y \gg 1$ (see Fig.4a). Accordingly, the momentum transfers q_1 and q_2 are a small fraction of the longitudinal components, $\frac{|q_1|}{|p_2|}$ and $\frac{|q_2|}{|p_1|} \ll 1$. We can, therefore, write $q_1^2 = q_{1t}^2$ and $q_2^2 = q_{2t}^2$. Expanding these momenta with respect to p_1 and p_2 we have

$$q_{1\mu} = x_1 p_{1\mu} + \beta_1 p_{2\mu} + q_{1t;\mu} , \quad (9)$$

$$q_{2\mu} = x_2 p_{2\mu} + \alpha_2 p_{1\mu} + q_{2t;\mu} .$$

We obtain

$$x_1 = \frac{\sqrt{M^2 + (q_1 - q_2)_t^2}}{\sqrt{s}} e^y \gg \alpha_2 , \quad (10)$$

$$x_2 = \frac{\sqrt{M^2 + (q_1 - q_2)_t^2}}{\sqrt{s}} e^{-y} \gg \beta_2 ,$$

where M denotes the Higgs mass.

We write the expression for the diagram of Fig.4a assuming that $k_t \gg q_{it}$ (see Fig.4a). Indeed, $q_{1t} \sim q_{2t} \approx \frac{1}{R}$, where R is the hadron (nucleon or nucleus) radius, while k_t can be as large as M . Inserting the vertex for Higgs emission by a gluon - gluon fusion [7], namely $\Gamma_H = g_H(k_{1\mu}k_{2\nu} - g_{\mu\nu}k_1.k_2)$, we obtain the following formula for the Higgs meson production amplitude of Fig.4a

$$A_H = \frac{1}{2}g_H \int dk_t^2 \phi(x_1, k_t^2) \phi(x_2, k_t^2) F_1(q_{1t}^2) F_2(q_{2t}^2) , \quad (11)$$

where $\alpha_S(k^2)xG(x, k^2) = \int^{k^2} dk'^2 \alpha_S(k'^2)\phi(x, k'^2)$, F_1 and F_2 are the form factors describing the t dependence of the Pomeron - target vertex ($F_1(0) = F_2(0) = 1$). We wish to stress that $\phi(x, k^2)$ are not the usual gluon densities, but the asymmetric (off diagonal) density functions that have been introduced in Ref.[8], and have been discussed in more detail in Ref.[9]. In the leading $\log(1/x)$ approximation of perturbative QCD [9], they can be approximated in the region of very small x_1 and x_2 by the usual nucleon gluon densities.

Neglecting the k^2 dependence of $\alpha_S(k^2)$, $\phi = \frac{dxG(x, k^2)}{dk^2} \propto (k^2)^{<\gamma>-1}$, where $<\gamma>$, the average anomalous dimension, can be considered to be a smooth function of both x and k^2 in the region of small x . This is a semi-classical approximation, which holds at small x [8], and we can use it to estimate the integral in question. Substituting ϕ in Eq. (11) we see that for all $<\gamma> < \frac{1}{2}$ the integral is infrared divergent. Only large distances contribute to the integral and we can, therefore, replace this integral by the exchange of a soft Pomeron as was done in Ref.[6]. In most of the models for the soft Pomeron [10], the real part of the Pomeron amplitude is very small, therefore, we expect that the new mechanism will only have a very small contribution to soft inclusive production.

In general, $<\gamma>$ is a function of x and k^2 , approaching the value $<\gamma> = \frac{1}{2}$ in some kinematic region. In the vicinity of $<\gamma> = \frac{1}{2}$ the integral over k^2 becomes logarithmic and all distances ($r_\perp$), down to very small $r_\perp \approx \frac{1}{M}$, contribute to the integral. In this case we need to consider two new physical phenomena: the BFKL Pomeron [11] and shadowing corrections (SC) [8] [12]. For the BFKL Pomeron, the value of the anomalous dimension cannot be larger than $\gamma = \frac{1}{2}$ so we evaluate our integral considering $\gamma \rightarrow \frac{1}{2}$. In this case

$$ReA_P = \frac{\pi}{2} \frac{dImA_P}{d\ln(1/x)} = \frac{\pi}{2} \omega_L ImA_P = 2\alpha_S N_c \ln 2 , \quad (12)$$

where $\alpha_P(0) = 1 + \omega_P$ and we have used the explicit calculation of $\omega_P = \omega_L$ [11]. One can see that for $\alpha_P(0) = 1.25$, $ReA_P \approx ImA_P$. Therefore, we expect a considerable violation of the factorization theorem for the BFKL Pomeron, as well as for a DGLAP Pomeron with an intercept which is well above unity [13].

An estimate of SC in the case of Higgs production via double soft Pomeron exchange in nucleon-nucleon scattering has been discussed in [12]. For the DGLAP evolution SC become important in the vicinity of $\gamma = \frac{1}{2}$, where $ImA_P \propto (\frac{1}{x})^{\omega_{cr}}$ [8] with $\omega_{cr} = \frac{2N_c\alpha_s}{\pi}$. Once again, for $\alpha_P(0) = 1.25$, we have $ReA_P \approx ImA_P$.

We can, thus, expect a considerable violation of the factorization theorem for hard inclusive production, in the kinematic region where the anomalous dimension approaches $\frac{1}{2}$ and where ω_p is well above 0.

To give a numerical estimate for the additional cross section we write

$$\frac{d\sigma^{NF}}{dy} = \int \frac{dq_{1t}^2 dq_{2t}^2}{(16\pi)^2} |A_H|^2 = \frac{2\pi^2}{R_1^2 R_2^2} \sigma(hard) \left| \int_{Q_0^2}^{M^2} dk_t^2 \frac{\partial \phi_1(x_1, k^2)}{\partial \ln(1/x_1)} \phi_2(x_2, k^2) \right|^2, \quad (13)$$

where $\sigma(hard)$ denotes the hard cross section, which is the same as in the factorization theorem (see Eq. (1)), R_1 and R_2 are the radii of the colliding hadrons (nuclei) normalized so that: $F_i(t) = 1 + \frac{1}{4}R_i^2 t$. For the nucleus - nucleus collision $\phi_i = A_i \phi_N$ and $R_i^2 = r_0^2 A_i^{\frac{2}{3}}$ and we get $\sigma^{NF} \propto A_1^{\frac{4}{3}} A_2^{\frac{4}{3}}$.

In Fig.5 we plot the ratio of the nonfactorized contribution of Eq. (13) to the cross section from the factorization theorem of Eq. (1), namely

$$R = \frac{2\pi^2}{R_1^2 R_2^2} \frac{\left| \int_{Q_0^2}^{\frac{M^2}{4}} dk_t^2 \frac{\partial \phi_1(x_1, k^2)}{\partial \ln(1/x_1)} \phi_2(x_2, k^2) \right|^2}{x_1 G_1(x_1, \frac{M^2}{4}) x_2 G_2(x_2, \frac{M^2}{4})}, \quad (14)$$

where $x_i G_i(x_i, M^2)$ is the gluon density of the i -th hadron. In Fig.5 we took $y = 0$. Notice, that this ratio does not depend on the hard cross section, and as such is also applicable to any central production process, in particular, hard minijet and jet production. In evaluating Eq. (14) we use the GRV parametrization [13] for the gluon density. We assume that $R_1^2 = R_2^2 = 5 \text{ GeV}^{-2}$ (see Ref.[14]). For the initial virtuality we take $Q_0^2 = 1 \text{ GeV}^2$, as the GRV parametrization is in agreement with the HERA data on $F_2(x, Q^2)$ [15] for all $Q^2 \geq 1 \text{ GeV}^2$.

One can see, that R is quite big for low masses and decreases when M increases. For example, $R = 1.2$ for $M = 10 \text{ GeV}$. We do not expect the Higgs meson mass to be that small, but our result is also applicable in the case of jet production (see Fig.4b) where $M \approx 2p_\perp$. Accordingly, with such a value of R we can expect minijet production (jets with $p_\perp \approx 5 \text{ GeV}$) which can be responsible for the structure of the minimum bias events at the Tevatron energy.

The value of R is bigger for nucleus - nucleus interaction. To estimate this value we need to multiply Eq. (14) by a factor of $A_{1eff}^{\frac{1}{3}} A_{2eff}^{\frac{1}{3}}$. Using the simple relation $R_A^2 = r_0^1 A^{\frac{2}{3}}$,

one has $A_{eff}^{\frac{1}{3}} = \frac{R_1^2}{r_0^2} (A^{\frac{1}{3}} - 1) + 1$. Therefore, for a gold - gold interaction we expect that R is enhanced by factor 4 - 9.

In calculating R we have not included a Sudakov form factor, which is believed to be necessary in a diagram of the type of Fig.5a, i.e. there is no emission of gluons with rapidities between y_1 and y_2 in Fig.4 (see Ref.[16] for details). However, as we have calculated an inclusive production process, this type of emission should be included (see below). As a first approximation we calculate the ratio R without the Sudakov form factor and neglect the BFKL enhancement, which can be important for large values of M (see Fig.6).

The diagrams of Fig.4 are only a low order approximation for the new process. Many additional gluons can be emitted, which lead to more complicated diagrams, shown in Fig.6. The emission of the additional gluons can be summed, and the cross section can be described by a Mueller diagram with the Pomeron - Pomeron interaction (λ) as seen in Fig.6. It is important to note that we get the same equation (see Eq. (8)) for the inclusive Higgs cross section, when using the AGK cutting rules for the Pomeron - Pomeron amplitude (λ) as was obtained in Ref.[17]. Indeed, it was shown in Ref.[17] that for different cuts of λ we have a transition matrix which can be written in the following form

$$\lambda_0^2 : \lambda_1^1 : \lambda_2^0 : \lambda_2^2 = 1 : 1 : \frac{1}{2} : \frac{1}{2} \quad (15)$$

with all other λ_i^f are equal to zero. Here, we denote by λ_i^f the elements of the transition matrix for the Pomeron - Pomeron amplitude, where i (f) is the initial (final) multiplicity state (say, above (below) the Pomeron - Pomeron interaction in Fig.6) and i (f) can be equal to 0,1 and 2, respectively. Using this transition matrix together with the AGK cutting rules for the inelastic cross section, one obtains the inclusive cross section for Higgs production

$$\begin{aligned} & \frac{d\sigma}{dy}(D + D \rightarrow H + X) \\ &= \rho^2 \frac{d\sigma}{dy}(D + D \rightarrow M_1(n \approx 1) + (LRG) + (H + X) + (LRG) + M_2(n \approx 1)) , \end{aligned} \quad (16)$$

where M_1 and M_2 denote the bunches of final particles with low multiplicities. Eq. (16) provides a method for calculating the inclusive Higgs production in terms of the cross section for the process with two large rapidity gaps, and the Higgs meson in the central rapidity region accompanied by a bunch of final hadrons.

The main result of our paper is Eq. (14), which gives the scale for the violation of the factorization theorem due to the new mechanism present in the inclusive production. It is clear, that this formula describes not only the process of a heavy Higgs meson production,

but also other central hard processes. The only difference is in the explicit expression for the hard cross section in Eq. (13). From Eq. (14) one can see that the new mechanism only gives a small contribution to the inclusive cross section for the production of light hadrons with small transverse momentum (soft inclusive production), because the real part of the elastic amplitude is very small at high energy. For hard inclusive production, the new mechanism yields a sizeable contribution, especially in the kinematic region where the anomalous dimension of the gluon density is close to $\frac{1}{2}$.

Our general conclusion is that the factorization theorem[1] is only approximate. We suggest that its proof should be reexamined with the view of finding the source of the violation.

References

- [1] J.C. Collins, D.E. Soper and G. Sterman: *Nucl. Phys.* **B308** (1988) 833.
- [2] A.H. Mueller: *Phys. Rev* **D23** (1970) 2963.
- [3] V.A. Abramovski, V.N. Gribov and O.V. Kancheli: *Sov. J. Nucl. Phys.* **18** (1973) 308.
- [4] M. Gyulassy, S.K. Kauffmann and L.W. Wilson: *Phys. Rev.* **C20** (1979) 2267;
I.V. Andreev, M. Plumer and R.M. Weiner: *Int. J. Mod. Phys.* **A8** (1993) 4577.
- [5] M.G. Ryskin: *Phys. Lett.* **B385** (1996) 365.
- [6] A. Bialas and P.V. Landshoff: *Phys. Lett.* **B256** (1991) 540.
- [7] T. Rizzo: *Phys. Rev.* **D 22** (1980) 178; S. Dawson and Russel P. Kauffmann: *Phys. Rev.* **D40** (1993) 2298 and reference therein.
- [8] L.V. Gribov, E.M. Levin and M.G. Ryskin: *Phys. Rep.* **100** (1983) 1.
- [9] M.G. Ryskin, R.G. Roberts, A.D. Martin and E.M. Levin: DTP/96/50, hep - ph/9606443, *Z. Phys.* **C** (in press); A.V. Radyushkin: CEBAF-TH-96-06, hep-ph/9605431; J.C. Collins, L. Frankfurt and M. Strikman: CERN-TH/96-314, hep-ph/9611433.
- [10] A. Donnachie and P.V. Landshoff: *Phys. Lett.* **B185** (1987) 403; *Nucl. Phys.* **B311** (1989) 509; E. Gotsman, E. Levin and U. Maor: *Z. Phys.* **C57** (1993) 677.

- [11] E.A. Kuraev, L.N. Lipatov and V.S. Fadin: *Sov. Phys. JETP* **45** (1977) 199 ; Ya.Ya. Balitskii and L.V. Lipatov: *Sov. J. Nucl. Phys.* **28** (1978) 822; L.N. Lipatov: *Sov. Phys. JETP* **63** (1986) 904.
- [12] E. Gotsman, E.M. Levin and U. Maor: *Phys. Lett.* **B353** (1995) 526.
- [13] M. Gluck, E. Reya and A. Vogt: *Z. Phys.* **C53** (1992) 127.
- [14] E. Gotsman, E. Levin and U. Maor: TAUP 2405/97, hep-ph/9701236.
- [15] H. Abramowicz: rapporteur talk in “*Proceedings of XXVIII ICHEP*”, Warsaw, July 25 - 27, 1996, TAUP 2396 - 96, hep-ph/9612001 and references therein.
- [16] A. Khoze, A.D. Martin and M.G. Ryskin: DTP-97-12, hep-ph/9701419.
- [17] J. Bartels and M.G. Ryskin: DESY 96-238, hep-ph/9612226.

Figure Captions

Fig.1: Mueller diagrams for inclusive Higgs production in a deuteron - deuteron collision :
 (a) the factorization theorem contribution, (b) the two nucleon interactions,
 (c) the diagrams, that cancel in total inclusive cross section, (d) the interference
 diagram that should be added to factorization theorem contribution.

Fig.2: The AGK cutting rules for inelastic and inclusive nucleon - nucleon cross sections.

Fig.3: Mueller diagrams for the new mechanism of the inclusive production:
 (a) $\sigma_2^{(0)}$, (b) $\sigma_2^{(1)}$, (c) $\sigma_2^{(2)}$ and the relation between different
 Mueller amplitudes ((d) and (e)).

Fig.4: Feynman diagrams (a) for Higgs and (b) for two high p_T jet production.

Fig.5: Calculated R (see Eq.(14)).

Fig.6: The general Mueller diagram for the new mechanism
 for the hard inclusive production.





Fig 1

$$\begin{aligned}
 \sigma_{in} &= \left| \begin{array}{c} \text{---} \\ | \text{ } \text{ } \text{ } | \\ \text{---} \end{array} \right|^2 - \left| \begin{array}{c} \text{---} \\ | \text{ } \text{ } | \\ \text{---} \end{array} \right|^2 = \\
 &= \left| \begin{array}{c} \text{---} \\ | \text{ } \text{ } | \\ \text{---} \end{array} \right|^2 + \left| \begin{array}{c} \text{---} \\ | \text{ } \text{ } | \\ \text{---} \end{array} \right|^2 - 2 \left| \begin{array}{c} \text{---} \\ | \text{ } \text{ } | \\ \text{---} \end{array} \right|^2 + \left| \begin{array}{c} \text{---} \\ | \text{ } \text{ } | \\ \text{---} \end{array} \right|^2 = \\
 &\quad \sigma^{(0)} + \sigma^{(1)} + \sigma^{(2)}
 \end{aligned}$$

$$\begin{aligned}
 \sigma_{incl} &= \left| \begin{array}{c} \text{---} \\ | \text{ } \text{ } | \\ \text{---} \end{array} \right|^2 - 2 \left| \begin{array}{c} \text{---} \\ | \text{ } \text{ } | \\ \text{---} \end{array} \right|^2 + 2 \left| \begin{array}{c} \text{---} \\ | \text{ } \text{ } | \\ \text{---} \end{array} \right|^2 = \\
 &= \left| \begin{array}{c} \text{---} \\ | \text{ } \text{ } | \\ \text{---} \end{array} \right|^2 = \left| \begin{array}{c} \text{---} \\ | \text{ } \text{ } | \\ \text{---} \end{array} \right|^2
 \end{aligned}$$

Fig. 2








Fig. 4









Fig 6



a)



b)